

Chandra and *XMM-Newton* view of the Warm Absorbing Gas in Mrk 290

Shuinai Zhang^{1*}, L. Ji², H. L. Marshall², A. L. Longinotti², D. Evans², Q. S. Gu^{1,3}

¹*Department of Astronomy, Nanjing University, Nanjing 210093, China*

²*MIT Kavli Institute, Cambridge, MA 02139, U.S.A.*

³*Key Laboratory of Modern Astronomy and Astrophysics (Nanjing University), Ministry of Education, China*

Accepted ??; Received ??; in original form ??

ABSTRACT

We present a detailed analysis of the *Chandra* High Energy Transmission Grating Spectrometer (HETGS) and *XMM-Newton* high resolution spectra of the bright Seyfert 1 galaxy, Mrk 290.

The *Chandra* HETGS spectra reveal complex absorption features that can be best described by a combination of three ionized absorbers. The outflow velocities of these warm absorbers are about 450 km s^{-1} , consistent with the three absorption components found in a previous far UV study. The ionizing continuum of Mrk 290 fluctuated by a factor of 1.4 during *Chandra* observations on a time scale of 17 days. Using the response in opacity of the three absorbers to this fluctuation, we put a lower limit on the distance from the ionizing source of 0.9 pc for the medium ionized absorber and an upper limit on distance of 2.5 pc for the lowest ionized absorber. The three ionization components lie on the stable branch of the thermal equilibrium curve, indicating roughly the same gas pressure. Therefore the thermal wind from the torus is most likely the origin of warm absorbing gas in Mrk 290.

During the *XMM-Newton* observation, the ionizing luminosity was 50% lower compared to the one in the *Chandra* observation. The RGS spectrum is well fitted by a two phase warm absorber, with several additional absorption lines attributed to a Galactic high velocity cloud, Complex C. Neither the ionization parameter ξ nor the column density N_{H} of the two absorbing components varied significantly, compared to the results from *Chandra* observations. The outflow velocities of both components were 1260 km s^{-1} . We suggest that an entirely new warm absorber from the torus passed through our line of sight.

Assuming the torus wind model, the estimated mass outflow rate is $\sim 1M_{\odot}$ per year, while the nuclear accretion rate is $\sim 0.5M_{\odot}$ per year. The O VII and Ne IX forbidden lines are the most prominent soft X-ray emission lines, with a mean redshift of 700 km s^{-1} relative to the systematic velocity. There seems to be no relation between emission lines and warm absorbers.

Key words: galaxies: Seyfert — galaxies: absorption lines — X-rays: galaxies — galaxies: individual: Mrk 290

* E-mail: snzhang@nju.edu.cn

1 INTRODUCTION

Ionized outflowing gas (or ‘warm absorber’, WA) is believed to be common in the inner parts of Active Galactic Nuclei (AGN) based on X-ray spectroscopy. About 50% of Seyfert 1s (Reynolds 1997; George et al. 1998) and quasars (Piconcelli et al. 2005; Misawa et al. 2007; Ganguly & Brotherton 2008) are characterized by blue-shifted X-ray absorption lines, the velocities of which are usually several hundred km s^{-1} . UV narrow absorption lines from outflowing gas were also detected at a similar fraction (Crenshaw et al. 1999), and showed velocities and ionization states as similar to WAs in some cases (Gabel et al. 2005a,b; Costantini et al. 2007). However, in other cases there is no clear physical correspondence between them (Kriss 2004).

Several studies have shown that WAs can be well modeled by including multiple absorbing components, which appear to be in pressure equilibrium (e.g. Netzer et al. 2003; McKernan et al. 2007; Krongold et al. 2009). Thus, the structure of WAs can be interpreted as a multi-phase medium, and most likely occupies the same spatial region (Krolik & Kriss 2001). Another suggestion that WAs occur in continuous density-stratified streamers (Behar et al. 2003; Steenbrugge et al. 2005) seems to be ruled out (Krongold et al. 2005, 2007).

Unfortunately, very little is still known about the physical states of WAs, or their dynamics and geometry. The most fundamental question is where WAs originate. Proposed locations span a wide range in radial distance from the central ionizing source: the accretion disk (Elvis 2000; Krongold et al. 2007), the broad line region (BLR) (Kraemer et al. 2005), the putative obscuring torus (Krolik & Kriss 2001; Blustin et al. 2005) and the narrow line region (Behar et al. 2003; Crenshaw et al. 2009). Simulations have offered similar possibilities, such as the accretion disk wind (Proga & Kallman 2004; Risaliti & Elvis 2009), the wind from torus (Dorodnitsyn et al. 2008), the large-scale outflows (Kurosawa & Proga 2009), but ruled out few options.

In principle, these outflows could potentially provide a common form of AGN feedback. But whether the outflow winds could extend into the interstellar medium and intergalactic medium (Nicastro et al. 2008) or just fall back into the accretion disk (Stoll et al. 2009), depends critically on the exact location of WAs when estimating the mass outflow rate and the total outflow kinetic energy. One key diagnostic for the location of WAs is based on applying non-equilibrium models on time variable AGNs (Nicastro et al. 1999). Such strong variability needed to apply to this method is not that common.

Mrk 290, a relatively nearby Seyfert 1 galaxy ($z = 0.0304 \pm 0.0014$ from the Sloan Digital Sky Survey), has been extensively studied by low resolution instruments (e.g. Wood et al. 1984; Kruper et al. 1990). The WA was found in the *ASCA* observation (Turner et al. 1996). The source flux of 2-10 keV band was $8.4 \times 10^{-11} \text{ erg s}^{-1} \text{ cm}^{-2}$ in the *ASCA* observation. In 2003, the first high resolution observation was carried out using *Chandra* HETGS, simultaneously with Far Ultraviolet Spectroscopic Explorer (*FUSE*) to determine the ionization state and kinematics of the WA, and assess association with the UV-absorbing gas. Mrk 290 was observed by *XMM-Newton* in 2006.

We present the analysis of the ionized absorbers in Mrk 290, using the high resolution X-ray spectra. This paper is organized as follows. In section 2, we describe the data reduction. Section 3 is devoted to the analysis of the data, and section 4 to the detailed model of the ionized absorbers of Mrk 290. In section 5 we discuss our results and present our conclusions in section 6. The cosmological parameters used are: $H_0 = 73 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_\Lambda = 0.72$ and $\Omega_m = 0.24$ (Spergel et al. 2007). The luminosity distance is 123.7 Mpc in this cosmological model. The quoted errors refer to 90% confidence for one interesting parameter.

2 OBSERVATION AND DATA REDUCTION

2.1 *Chandra* Data: 2 Flux States

Chandra performed four observations of Mrk 290 for a total duration of 251 ks, using the High Energy Transmission Grating Spectrometer (HETGS, Canizares et al. 2005). Table 1 lists the observation log.

The HETGS is composing of two grating sets, the High Energy Gratings (HEG) and the Medium Energy Gratings (MEG), the resolutions of which are 0.012 Å and 0.023 Å, respectively. All observations were reduced uniformly and in a standard way using the *Chandra* Interactive Analysis of Observations (CIAO) software (Version 4.1.1). Flux calibration of each observation was carried out using the *Chandra* Calibration Database (Version 4.1.2). The +1st and -1st orders of each HEG and MEG spectra were combined; higher orders were ignored.

We generated the light curves in 1000 second bins for the short wavelength band (2-10 Å), as shown in Fig. 1. The comparison between them indicates that the time-average count rate increased by a factor of ~ 2 , between observations 4399 and 3567, only half a month apart. Smaller variations are observed on time scales of 10^4 s within each observation, which were also found by Turner et al. (1996). Figure 1 shows the softness ratio, defined as the count rate in the 15-25 Å band divided by that in the 2-10 Å band, to illustrate the different variability characteristics below and above 1 keV (Netzer et al. 2003). The softness ratio increases from 0.075 to 0.100 along with a higher count rate, between observations 4399 and 3567. In light of these results, we have divided the entire dataset into two groups with higher (ID 3567, 4441 & 4442) and lower (ID 4399) count rates.

2.2 *XMM-Newton* Data Reduction

Mrk 290 was observed four times by *XMM-Newton* (Jansen et al. 2001) during the period April 30 to May 6, 2006. The European Photon Imaging Camera (EPIC) was operated in small window mode. Both RGS (Reflection Grating Spectrometer) cameras were in spectroscopic

mode. Raw data from all observations were processed with the standard Science Analysis System (SAS) using version 8.0 and the most updated calibration files.

High ‘flaring’ background periods were all filtered to maximize the signal-to-noise ratio (S/N). While the total exposure time is 100 ks, after this screening process, the net exposure time was 65.5 ks, as shown in Table 2. The four *XMM-Newton* data were merged, since no significant flux variability was found.

The *pn* spectra were extracted using pattern 0-4. The 0.5-10 keV count rate, $2.5 \text{ counts s}^{-1}$, was under the pile-up threshold, verified with the EPATPLOT. Each source spectrum was extracted from a circular region of $40''$ of radius centered on the peak of the X-ray emission. The background spectrum was taken from a vicinity source-free circular region of radius of $55''$. RGS data were processed with the task RGSPROC according to the standard method proposed in the SAS thread.

3 SPECTRAL ANALYSIS

We used the Interactive Spectral Interpretation System (ISIS version 1.5.0, (Houck 2002)) to fit the spectra. When fitting the data, we used the *C* statistic (Cash 1976) to find the best-fitting model parameters in the high-resolution spectra.

3.1 The Broadband Continuum Modeling

The Mrk 290 spectrum is affected by a relatively low Galactic absorption: $N_H = 1.76 \times 10^{20} \text{ cm}^{-2}$ (Kalberla et al. 2005), which is included in all models.

The *Chandra* spectra were binned to 0.01 Å , and fitted in the wavelength range 1.5-25 Å, as shown in Fig. 2. In contrast to previous results, where Mrk 290 was described by an absorbed power law (Turner et al. 1996; Shinozaki et al. 2006; Jiang et al. 2006), we found that a black-body component was needed in addition to the absorbed power law in order to fit the continuum. The black-body component was included to account for the soft excess, that could come from a standard optically-thick accretion disk. The intrinsic cold absorption was negligible compared to the Galactic column density. In fact, half of AGN (McKernan et al. 2007) and quasars (Piconcelli et al. 2005) do not appear to have significant intrinsic cold gas absorption.

The *pn* spectrum was grouped to give a minimum of 50 counts per bin in order to apply χ^2 statistics. The same model as fitted to HETGS spectra but with free parameters provided a good fit to the *pn* data, except for two prominent residuals around 0.8 and 6.3 keV. These two features are consistent with typical absorbing features of WA and Fe $\text{K}\alpha$ emission line, respectively. Again, no cold absorption component above Galactic was needed. Fig. 3 shows the spectrum in the 0.5-10 keV band.

The best fit parameters of all spectra are given in Table 3, including the fluxes and luminosities in the 2-10 keV band. The hard X-ray luminosity increased by a factor of 1.4 in 17 days between *Chandra* observations. When *XMM-Newton* re-observed the source three years later, the luminosity showed a decrease of 50% compared to the highest value measured in the *Chandra* spectra. So hereafter, we refer to these spectra as ‘high-’, ‘mid-’ and ‘low-state’ spectra as labeled in Table 3.

3.2 Intrinsic Absorption Line Properties

The HETGS spectra, which are binned to 0.01 Å , reveal many absorption line features, some of which are blends of several lines. H-like and He-like lines of O, Ne, Mg, Si, S, and L-shell lines of Si, Fe are marked in Fig. 2. The M-shell complex of iron, called the unresolved transition array (UTA) is observed as a depression at 16-17.5 Å, although it is very shallow in the present data. We identified one prominent narrow absorption line of Fe x in the UTA (see Table 4). All statistically significant lines with Poisson probability greater than 0.995 were measured by fitting a negative Gaussian to the continua within $\pm 0.5 \text{ Å}$ band, as shown in Table 4. The widths of the Gaussian lines were all fixed to 1 mÅ . The outflow velocities about 450 km s^{-1} were derived with respect to the systemic redshift 0.0304, except those lines labeled as Galactic absorption. There are fewer significant lines in the mid-state spectrum than in the high-state spectrum.

The RGS spectra are binned to 0.1 Å , in which the iron UTA shows clearly. We fitted RGS 1 and 2 simultaneously in the wavelength band 7-35 Å, but for clarity added them in Fig. 4. The continuum model was fixed to that obtained from *pn* spectrum, except for allowing the black body component to vary. We found several highly ionized features: Ne ix combined with Fe xix, Ne x Ly α , C vi, N vii, Fe xviii and Fe xx complex. A Gaussian was fitted to each feature and the results were listed in Table 4. The outflow velocities of these lines are more than 1000 km s^{-1} , differing by the several hundred km s^{-1} from those derived from the HETGS spectra.

3.3 Galactic Absorption

The spectra of Mrk 290 show narrow absorption lines at zero redshift in Table 4. O i absorption at 23.5 Å in Fig. 4 is attributed to the ISM of our Galaxy (Wilms et al. 2000). We also detected ionized material traced by O viii in the HETGS spectra, and O vii, N vii, N vi, and possibly C v in the RGS spectrum. This ionized absorber may be associated with the high-velocity cloud complex C, which is a low-metallicity gas cloud plunging toward the disk and beginning to interact with the ambient gas that surrounds the Milky Way (Collins et al. 2007). In the UV band, Penton et al. (2000) detected absorption lines from complex C using HST observations. Since this ionized gas does

not affect the continuum shape (Costantini et al. 2007), we fitted these lines using Gaussians. The upper limits of the infall velocity with respect to our Milky Way were measured from these lines less than 1000 km s^{-1} , consistent with the values from the UV lines: 200 km s^{-1} and 120 km s^{-1} (Collins et al. 2007).

3.4 Emission Lines

In the soft X-ray band, Mrk 290 exhibits a prominent Ne IX forbidden line in the high-state spectrum and a O VII forbidden line in the high- and low-state spectra. They are red-shifted at offset velocities of 700 km s^{-1} with respect to the systemic redshift, which is definitely different from the absorption lines. The velocities do not change over three years suggests that these emission lines may be emitted in the same gas. The fit results for the three lines are shown in Table 5.

The iron emission line at 6.4 keV (or $1.9378 \text{ \AA}$) is present in both the high-state HETGS spectrum and the low-state *pn* spectrum. This fluorescent line is $K\alpha$ transitions from neutral iron, consistent with Fe I-XII. A Gaussian line, the width of which was fixed at $1 \text{ m\AA}$, had been included to fit the Fe $K\alpha$ line in the $5\text{--}8 \text{ keV}$ band. Table 5 lists the measurements. The equivalent widths (EWs) of $48 \pm 17 \text{ eV}$ for the high-state spectrum and of $23 \pm 17 \text{ eV}$ for the low-state spectrum are consistent with the result of Shu et al. (2010). However they are much smaller than the EWs from ASCA data: 273 eV (Shinozaki et al. 2006) or 500 eV (Turner et al. 1996).

We allowed the line width to float and fitted the iron $K\alpha$ again. The Cash statistics changed by 2 and 8 when reducing one *d.o.f.* for the high- and low-state spectra, respectively, and the equivalent widths (EWs) increased to $71 \pm 35 \text{ eV}$ and $81 \pm 29 \text{ eV}$. The FWHM of Fe $K\alpha$ in the high-state spectrum is $5700^{+9100}_{-4600} \text{ km s}^{-1}$, and in the low-state *pn* spectrum is $14000^{+10000}_{-6000} \text{ km s}^{-1}$, which agrees with ASCA result (Turner et al. 1996).

In the high-state spectrum, a line at $1.897 \text{ \AA}$ seems consistent with the $n=2\text{--}1$ resonance transition of Fe XXV (at a rest energy of 6.696 keV). This line could be produced in Compton-thin, low velocity photoionized material (Bianchi et al. 2005).

There are unidentified emission features with Poisson probabilities less than 0.001 in the high-state spectrum at $2.666 \text{ \AA}$ and $8.725 \text{ \AA}$, of which the fluxes are about $few \times 10^6 \text{ ph cm}^{-2} \text{ s}^{-1}$. However, this two features are probably spurious.

4 PHOTOIONIZATION MODELING

The publicly available photoionization code XSTAR¹ is the tool we used to model the physical conditions of the absorbing gas. It includes H- and He-like transitions of most ions, L-shell transitions of Fe, and important M-shell transitions of Fe that contribute to the UTA feature. We used PVM_XSTAR (Noble et al. 2009) to generate grids of models, which provides parallel execution of XSTAR 2.1ln11.

We adopted the default solar abundances (Grevesse et al. 1996) in XSTAR, and used a velocity turbulence (*b*-value) of 170 km s^{-1} , which corresponds approximately to the limiting MEG spectral resolution ($\sim 300 \text{ km s}^{-1}$ FWHM at 0.5 keV) (McKernan et al. 2007). This choice is justified by the absence of detectable line broadening.

The photoionization model used here contains three parameters: red-shift *z*; total neutral hydrogen column density N_H ; and the ionization parameter $\xi = L_{ion}/(n_e R^2)$, where L_{ion} is the ionizing luminosity in the range $1\text{--}1000 \text{ Ryd}$, n_e is the electron density, and R is the distance of the ionized gas from the central ionizing source.

4.1 The Spectral Energy Distribution

First of all, we constructed the spectral energy distributions (SEDs) of Mrk 290 to generate the photoionization models. However, the intrinsic X-ray spectrum, an important part of the SED, is obtained by fitting the X-ray data with photoionization models. Thus a two-step iterative process is employed to set the intrinsic spectrum, as described in Yaqoob et al. (2003).

In the first step, the ionizing X-ray spectrum is obtained by fitting the continuum without a WA to construct a SED. The SED is then used to generate a grid of warm absorption models that are fit to the spectra. The second step is similar to the first step: using the new intrinsic X-ray spectrum for a new SED, which is in turn used to generate new set of photoionization models.

The *Chandra* mid-state data (ID 4399) was observed simultaneously in the UV band by *FUSE*. We adopted this UV data point (Dunn et al. 2008), while other data were obtained from NASA/IPAC Extragalactic Database (NED): radio data from VLA, 25μ data from IRAS, and r band data from SDSS.

For each state, the flux point at 0.5 keV band on the intrinsic X-ray continuum was simply linked to the UV point and to other data in the SED by straight lines in log-log space, as shown in Fig 5. We derived luminosities of $0.5\text{--}13.6 \text{ keV}$ band from the intrinsic X-ray spectral models, measured luminosities of $13.6 \text{ eV} - 0.5 \text{ keV}$ band from the two point lines in SEDs, and added them to determine L_{ion} . Detailed intrinsic continuum parameters and luminosities are listed in Table 7.

¹ <http://heasarc.gsfc.nasa.gov/docs/software/xstar/xstar.html>

4.2 Warm Absorbers from Model Results

4.2.1 One WA Component

Turner et al. (1996) modeled the *ASCA* spectrum of Mrk 290 with one WA component generated using XSTAR, obtaining an ionization parameter $\log(\xi) = 1.38$ and a column density N_H of $8 \times 10^{21} \text{ cm}^{-2}$. Similarly, we began with one WA in fitting *Chandra* HETGS spectra. The *Chandra* spectra were binned to $0.01 \text{ \AA}$, and fitted with one ionization component. The model represents most absorption lines well, and the best fit results are shown in Table 6. The absorber is blue-shifted by $-450 \pm 30 \text{ km s}^{-1}$ relative to the systemic redshift, and neither its column density nor ionization parameter changed apparently from mid-state to high-state during half a month. Since the UTA feature is quite clear in the RGS spectrum, one WA component fitting is not suitable for the low-state data.

4.2.2 Two Phase Warm Absorber

Coarsely binned high- and mid-state spectra are shown in Fig. 6, in which the green lines represent the best fitting model from Table 6. The visual inspection of the spectra reveals the unfitted UTA feature clearly in the $16\text{--}17.5 \text{ \AA}$ band, suggesting that the single WA component is not sufficient to account for all absorbing features.

The UTA is a broad feature, and its shape is sensitive to the charge states of Fe producing it, so we rebinned the spectra heavily. Since the UTA is a key diagnostic of material in a low-ionization state, a second WA with a lower ionization level is required to be included in the model. We modeled the high-state spectrum using a two phase WA with parameters set free, except the redshift of the higher ionization component was fixed to $z = 0.0289$ from Table 6. The two phase warm absorber provides a good fit to the high-state spectrum as shown in Fig. 6 by the red line model. The Fe UTA is well fitted by a lower ionized component with $\log(\xi) = 1.62 \pm 0.15$, $N_H = 5.4(\pm 1.8) \times 10^{20} \text{ cm}^{-2}$. The outflow velocity is similar to that of the high ionization component, and the other best fit parameters are reported in Table 7.

Providing that no properties in Table 7 varied in 17 days and the dynamical timescale is much longer than the radiative recombination timescale, when fitting the mid-state spectrum, we fixed the redshift and column density of each absorbing component to that of high-state, but allowed the ionization parameter to vary. Unfortunately, the large uncertainties in ξ of the low ionization component cannot constrain the possible response of opacity to the continuum variation.

The two phase WA model was also used to fit the RGS spectrum, adding several gaussian lines to account for the local absorption lines and the O VII forbidden emission line. Fig. 7 shows the best model for RGS spectrum, and fitting results are listed in Table 7. As expected, the deep UTA feature comes from the lower ionized material, while other features identified as transitions of Ne IX, Ne X, Fe XVIII, Fe XX and N VII, come from the higher ionization component. The C VI absorption lines are believed to be associated with the low ionization component (Sako et al. 2001), though the observed lines are much deeper than what the photoionization model predicts here. However, it could also be the result from carbon-rich gas. The Fe XVIII complex is not well fitted here, as in HETGS spectra (as shown in Fig. 6), which is partly due to the uncertainties of di-electronic recombination rates for iron L-shell ions (Gu 2003), and partly due to the uncertainty of the iron abundance (solar abundance is used in the photoionization models). Neither the ionization parameter nor the column density of the two WA components varied significantly compared to that found in the HETGS spectra, but the outflow velocity increased by a factor of about 2.5.

4.2.3 The Third, Hottest Ionization Component

We have applied the two phase warm absorber model back to the $0.01 \text{ \AA}$ binned high-state spectrum again and checked residuals carefully. Three important Fe L-shell absorption lines were left unfitted, Fe XXI ($12.284 \text{ \AA}$), Fe XXII ($11.770 \text{ \AA}$), Fe XXIII ($10.981 \text{ \AA}$). A hotter high ionization component is required to explain these absorption lines, as labeled in Fig. 8. Such a very high ionization absorbing component has been reported in a few sources, e.g. NGC 3783 (Netzer et al. 2003), NGC 5548 (Steenbrugge et al. 2005) and NGC 985 (Krongold et al. 2009).

We added the third ionized absorber in the high resolution high-state spectrum to test for the possible presence of this component. The lowest ionization component was fixed to the values of the two phase WA model fit, where they were best determined. The fit parameters are given in Table 8. The Cash statistic gives a difference of 45 for adding 3 d.o.f (2590/2341) compared to the model with one WA. The model of three ionization components are shown in Fig. 8. The comparison of the best-fitting model to the high-state spectrum can be seen in Fig. 9, in which the flux spectrum is binned to $0.025 \text{ \AA}$ for clarify.

4.3 The L-shell silicon lines

In *Chandra* HETGS spectra, Si L-shell absorption lines do not exist in the fitted model, because so far they have mostly been omitted from the XSTAR database (McKernan et al. 2007). Fig. 10 shows these unfitted lines in the mid- and high-state spectra. The mid-state spectrum shows Si XI, Si IX, and one significant line (at $7.23 \text{ \AA}$) that could be Si VII with an outflow velocity 2000 km s^{-1} , Si VIII with an inflow velocity 700 km s^{-1} , or spurious. In the high-state spectrum Si XI, Si X, and Si IX are evident.

The $7.23 \text{ \AA}$ line disappeared totally while Si IX got stronger from the mid-state to the high-state. Furthermore the ratio of Si IX to Si XI changed from 0.6 ± 0.5 to 1.6 ± 0.4 over the two states. Netzer (2004) suggested that the Si L-shell absorption lines should be generated from the same ionization component producing the Fe UTA. Gu et al. (2006) calculated the new Fe M-shell data and confirmed

this suggestion, especially for the NGC 3783. The variation of Si L-shell lines in Mrk 290 demonstrated the change of the ionization parameter in the lowest ionization component.

5 DISCUSSION

5.1 The Location of The WAs

When the ionizing luminosity L_{ion} and the ionization parameter ξ are known, one would expect to constrain the distance R by estimating the electron density n_e , since $\xi = L_{ion}/(n_e R^2)$. Studying the variation based on non-equilibrium models is one method to constrain n_e (Nicastro et al. 1999). An excellent approximation for the time to reach a new photoionization equilibrium is

$$t_{eq}^{x^i, x^{i+1}} \sim \left[\frac{1}{\alpha_{rec}(x^i, T_e)_{eq} n_e} \right] \times \left[\frac{1}{[\alpha_{rec}(x^{i-1}, T_e)/\alpha_{rec}(x^i, T_e)]_{eq} + [n_{x^{i+1}}/n_{x^i}]} \right], \quad (1)$$

where x_i is the relative density of ion i of the element X , and the recombination time $\alpha_{rec}(x^i, T_e)$ can be obtained from Shull & van Steenberg (1982) data.

During *Chandra* observations, the luminosity increased by a factor of 1.4, thus the $\log(\xi)$ was expected to increase 0.15 dex. However, the ionization parameter of the medium ionization component, which is best constrained in the *Chandra* spectra, did not change (Table 6). Providing the flux increase happened between June 29 and July 15 in 2003 and the high-state observation lasted for 2.5 days, the timescale for the gas to reach the new photoionization equilibrium must be longer than 2.5 days. Considering the ions O VI - O VIII and Ne VIII - Ne X, we find the electron density of the medium ionization component $n_e < 6.2 \times 10^4 \text{ cm}^{-3}$. Combined with L_{ion} and ξ , the distance from the center black hole to this WA is further than 0.9 pc.

The large error bars of ξ in the lowest ionization component make it hard to check whether ξ varied. However, based on the variation of Si L-shell lines, we conclude that the ionization state of the lowest ionization component changed over the 17 day span. Using the same method as above, we find that $n_e > 5.5 \times 10^4 \text{ cm}^{-3}$ and that the distance to this component is less than 2.5 pc. It might be common that a lower ionization component reaches a new photoionization equilibrium easier than a higher ionization components (Krongold et al. 2005), as in NGC 3783, NGC 4051, NGC 5548.

5.2 On The Origin of The Emission Lines

The location of gas producing soft X-ray emission lines is unclear. These lines may originate from just the WA (Netzer et al. 2003), very close to or within the BLR (Costantini et al. 2007; Longinotti et al. 2008, 2010), within the Narrow Line Region (NLR) (Guainazzi & Bianchi 2007), or between the BLR and the NLR (Nucita et al. 2010). The soft X-ray emission lines in Mrk 290 are redshifted with respect to the systemic velocity, and the velocity did not change over 3 years, so they are probably not related to the WAs.

The FWHM of Fe $K\alpha$ is similar to the width of H β (5323.5 km s⁻¹; Vestergaard & Peterson 2006), indicating a possible origin in the BLR (Bianchi et al. 2008). However, forcing the lines to be narrow does not make the fits significantly worse, which suggests that the torus is still the primary origin for this fluorescent line (Bianchi et al. 2007).

5.3 The Torus Origin

With the reverberation mapping results (Denney et al. 2010), the black hole mass of Mrk 290 is $2.43 \pm 0.37 \times 10^7 M_\odot$. The time delay of H β , $8.72^{+1.21}_{-1.02}$ days, indicates that the radius of the BLR is 0.007 pc. The thermal bolometric luminosity calculated from the SED, $L_{bol} = 2.5 \times 10^{44} \text{ erg s}^{-1}$, gives the radius of the torus: 1.1pc (e.g. Krolik & Kriss 2001). Thus, the radius of the torus seems similar to the distance of the WAs.

The thermal photoionization equilibrium curve (Krolik et al. 1981) is presented in Fig. 11 for the high-state SED. The curve marks the points of thermal equilibria in $T - \Xi$ plane, where T is the electron temperature and the “pressure” ionization parameter $\Xi = \xi/4\pi ckT$. The three WAs best modeled in the high-state spectrum are roughly in pressure balance as shown in Fig. 11, indicating that all three components may occupy the same spatial region. Combined with the distance, we suggest that the origin of the WAs is a thermal wind from the torus, the model proposed by Krolik & Kriss (2001).

The WAs in the RGS spectrum do not show significantly different ionization parameters and column densities compared to that found from the HETGS spectra. This supports the model that the WAs come from the torus. The large change of the outflow velocity, from 450 km s^{-1} to 1260 km s^{-1} , gives a large acceleration 0.6 cm s^{-2} . Radiation pressure is an attractive means for accelerating the gas (Crenshaw et al. 2003), and the acceleration can be written as $a = \frac{\kappa L}{4\pi r^2 c} - \frac{GM}{r}$, where κ is the absorption cross-section per unit mass. For Mrk 290, the acceleration by radiation pressure is $\sim 0.01 \text{ cm s}^{-2}$, much less than the observed one. As a result, an entirely new WA gas may have passed through our line of sight during the *XMM-Newton* observation.

A natural geometry for WA winds is bi-conical (Dorodnitsyn et al. 2008). Assuming this, the mass loss rate can be estimated. We use a formula for the mass outflow rate derived by Krongold et al. (2007):

$$\dot{M}_{out} = 0.8\pi m_p N_H v_r R f(\delta, \phi) \quad (2)$$

where $f(\delta, \phi)$ is a factor that depends on the particular orientation of the disk and the wind, and for all reasonable angles ($\delta > 20^\circ$ and $\phi > 45^\circ$) is of order of unity. With the observed v_r (the line of sight outflow velocity), N_H , and the distance of torus $R = 1.1$ pc, the mass outflow rate contributed by the medium and lowest ionized WAs is $\sim 1M_\odot$ per year, of the same order as the values in the simulation (Dorodnitsyn et al. 2008). The outflow mass rate estimate is larger than the accretion rate, $\dot{M}_{acc} = \frac{L_{bol}}{c^2 \eta} \sim 0.5M_\odot$ per year, assuming $\eta = 0.1$.

The escape velocity ($v_{esc} = \sqrt{2GM/R}$) from the location of the torus is about 440 km s^{-1} , which is nearly identical to the outflow velocities of the WAs in Mrk 290. The outflow velocity is only a lower limit to the actual speed, thus the warm gas would not likely fall back, and would move 5 kpc away from the central region into the host galaxy over the AGN lifetime (10^7 yr). The total mass lost due to the WAs is $M_{out} \sim 10^7 M_\odot$, and the integrated kinetic energy (given by $M_{out} v^2/2$) is $\sim 2 \times 10^{55}$ ergs in 10^7 yr. As suggested by Krongold et al. (2007), the hot interstellar medium in the host galaxy can be heated to 10^7 K and would evaporate, ceasing star formation. As a consequence, this outflow mass rate may affect the evolution of the host galaxy and the surrounding intergalactic medium, over the AGN lifetime.

5.4 The Connection between UV and X-ray Absorbers

Mrk 290 was observed by *FUSE* simultaneously with *Chandra*. Kaiser et al. (2006) detected three intrinsic absorbers by the O VI doublet in the UV band, blueshifted by 526, 470 and 439 km s^{-1} with respect to the systemic velocity. Comparing them with the outflow velocities of the three WAs in X-ray band, 540 ± 150 , 450 ± 30 and $390 \pm 60 \text{ km s}^{-1}$, we find the UV absorption components correspond to the X-ray WAs within the uncertainties. One geometry to interpret this connection is that high-density, low-ionization UV-absorbing clouds are embedded in a low-density, high-ionization material that dominates the X-ray warm absorber (Kriss 2004). Kaiser et al. (2006) estimated that the hydrogen column density is $9 \times 10^{14} \text{ cm}^{-2}$. Assuming $n_e > 6 \times 10^4 \text{ cm}^{-3}$, the thickness of the UV-absorbing clouds must be less than $1.5 \times 10^5 \text{ km}$. In fact, many X-ray observed absorption components appear to be accompanied by corresponding UV absorption components with the same outflow velocity (e.g. Steenbrugge et al. 2009).

6 CONCLUSIONS

We summarize the main results of this analysis of the Mrk 290 high resolution X-ray spectra as follows:

1. The *Chandra*-HETGS spectra reveal complex absorption features including the Fe M-shell unresolved transition array (UTA), indicating the presence of intervening ionized absorbing gas. Photoionization modeling shows that a combination of three ionization absorbers with similar kinematics (outflow velocities $\sim 450 \text{ km s}^{-1}$) can fit the data and are consistent with the three absorption components in the previous far UV band study. The ionized component with highest ionization is required by the presence of several absorption lines of Fe XXII-XXIII, which are rarely observed.

2. The ionizing continuum increased 40% during the *Chandra* observation, on a timescale of 17 days. Using the lack of response in the opacity of the higher ionization component to this fluctuation, we constrain the distance from the ionizing source to > 0.9 pc. Si L-shell line ratios show significant changes and are related to the UTA, so they result from same gas with the ionization level. If the variation derived from the Si L-shell lines of the lowest ionization component is real, we may set an upper limit of 2.5 pc for the distance to the central source. The three ionization components lie on the stable branch of the S-curve, showing roughly the same gas pressure, which indicates that they occupy the same spatial region. We conclude that the warm absorbing gas originates in the thermal wind from the torus.

3. During the *XMM-Newton* observation, the ionizing luminosity of Mrk 290 was only half of the high luminosity of the *Chandra* observation. The RGS spectrum is well fitted by a two phase warm absorber, with several additional local absorption lines attributed to the high velocity cloud Complex C. Neither the ionization parameter ξ nor the column density N_H of the two absorbing components varied significantly, compared to the results from *Chandra* observations. The outflow velocities of both components increased by a factor of 2.5. However, the radiation pressure in Mrk 290 is not enough to accelerate the gas to $\sim 1260 \text{ km s}^{-1}$. We suggest that entirely new WA gas from the torus passed through our line of sight.

4. The estimated mass outflow rate of the ionized wind is $\sim 1M_\odot$ per year, larger than the nuclear accretion rate $\sim 0.5M_\odot$ per year, and the input kinetic energy is $\sim 2 \times 10^{55}$ ergs.

5. In both high- and low-state spectra the Fe K α emission lines are detected. The O VII and Ne IX forbidden lines are the most prominent soft X-ray emission lines, with a mean redshift velocity of $\sim 700 \text{ km s}^{-1}$ with respect to the systematic velocity. There seems to be no relation between emission lines and warm absorbers.

ACKNOWLEDGEMENTS

We thank the help from the scientists of the MIT Kavli Institute, and Tim Kallman for detailed instructions about XSTAR. We thank the anonymous referee for providing useful comments on our work. Support for this work was provided by the National Aeronautic Space Administration through the Smithsonian Astrophysics of Observation contract SV3-73016 to MIT for support of the Chandra X-ray Center,

which is operated by SAO for and on behalf of NASA under contract NAS8-03060, by Program for New Century Excellent Talents in University (NCET), and by Chandra grant GO8-9113B. Also our work is supported under the National Natural Science Foundation of China under grants (10878010, 10221001 and 10633040) and the National Basic Research Program (973 programme no. 2007CB815405).

REFERENCES

- Behar, E., Rasmussen, A. P., Blustin, A. J., Sako, M., Kahn, S. M., Kaastra, J. S., Branduardi-Raymont, G., & Steenbrugge, K. C. 2003, *ApJ*, 598, 232
- Bianchi, S., Matt, G., Nicastro, F., Porquet, D., & Dubau, J. 2005, *MNRAS*, 357, 599
- Bianchi, S., Guainazzi, M., Matt, G., & Fonseca Bonilla, N. 2007, *A&A*, 467, L19
- Bianchi, S., La Franca, F., Matt, G., Guainazzi, M., Jimenez Bailón, E., Longinotti, A. L., Nicastro, F., & Pentericci, L. 2008, *MNRAS*, 389, L52
- Blustin, A. J., Page, M. J., Fuerst, S. V., Branduardi-Raymont, G., & Ashton, C. E. 2005, *A&A*, 431, 111
- Canizares, C. R., et al. 2005, *PASP*, 117, 1144
- Cash, W. 1976, *A&A*, 52, 307
- Collins, J. A., Shull, J. M., & Giroux, M. L. 2007, *ApJ*, 657, 271
- Costantini, E., et al. 2007, *A&A*, 461, 121
- Costantini, E., Kaastra, J. S., Korista, K., Ebrero, J., Arav, N., Kriss, G., & Steenbrugge, K. C. 2010, arXiv:1001.2712
- Crenshaw, D. M., Kraemer, S. B., Boggess, A., Maran, S. P., Mushotzky, R. F., & Wu, C.-C. 1999, *ApJ*, 516, 750
- Crenshaw, D. M., Kraemer, S. B., & George, I. M. 2003, *Annual Review of Astronomy & Astrophysics*, 41, 117
- Crenshaw, D. M., Kraemer, S. B., Schmitt, H. R., Jaffe, Y. L., Deo, R. P., Collins, N. R., & Fischer, T. C. 2009, arXiv:0912.2420
- Denney, K. D., et al. 2010, *IAU Symposium*, 267, 201
- Dorodnitsyn, A., Kallman, T., & Proga, D. 2008, *ApJ*, 687, 97
- Dunn, J. P., Crenshaw, D. M., Kraemer, S. B., & Trippe, M. L. 2008, *AJ*, 136, 1201
- Elvis, M. 2000, *ApJ*, 545, 63
- Gabel, J. R., et al. 2005a, *ApJ*, 631, 741
- Gabel, J. R., et al. 2005b, *ApJ*, 623, 85
- Ganguly, R., & Brotherton, M. S. 2008, *ApJ*, 672, 102
- George, I. M., Turner, T. J., Netzer, H., Nandra, K., Mushotzky, R. F., & Yaqoob, T. 1998, *ApJS*, 114, 73
- Grevesse, N., Noels, A., & Sauval, A. J. 1996, *Cosmic Abundances*, 99, 117
- Gu, M. F. 2003, *ApJ*, 590, 1131
- Gu, M. F., Holczer, T., Behar, E., & Kahn, S. M. 2006, *ApJ*, 641, 1227
- Guainazzi, M., & Bianchi, S. 2007, *MNRAS*, 374, 1290
- Houck, J. C. 2002, *High Resolution X-ray Spectroscopy with XMM-Newton and Chandra*
- Jansen, F., et al. 2001, *A&A*, 365, L1
- Jiang, P., Wang, J. X., & Wang, T. G. 2006, *ApJ*, 644, 725
- Kaiser, M. E., Lee, J. C., Kriss, G. A., Marshall, H., & Fang, T., Gibson, R. R. 2006, *Astrophysics in the Far Ultraviolet: Five Years of Discovery with FUSE*, 348, 522
- Kalberla, P. M. W., Burton, W. B., Hartmann, D., Arnal, E. M., Bajaja, E., Morras, R., Pöppel, W. G. L. 2005, *A&A*, 440, 775
- Kraemer, S. B., et al. 2005, *ApJ*, 633, 693
- Kriss, G. A. 2004, *The Interplay Among Black Holes, Stars and ISM in Galactic Nuclei*, 222, 223
- Krolik, J. H., McKee, C. F., & Tarter, C. B. 1981, *ApJ*, 249, 422
- Krolik, J. H., & Kriss, G. A. 2001, *ApJ*, 561, 684
- Krongold, Y., Nicastro, F., Brickhouse, N. S., Elvis, M., & Mathur, S. 2005, *ApJ*, 622, 842
- Krongold, Y., Nicastro, F., Elvis, M., Brickhouse, N., Binette, L., Mathur, S., & Jiménez-Bailón, E. 2007, *ApJ*, 659, 1022
- Krongold, Y., et al. 2009, *ApJ*, 690, 773
- Kruper, J. S., Canizares, C. R., & Urry, C. M. 1990, *ApJS*, 74, 347
- Kurosawa, R., & Proga, D. 2009, *MNRAS*, 397, 1791
- Longinotti, A. L., Nucita, A., Santos-Lleo, M., & Guainazzi, M. 2008, *A&A*, 484, 311
- Longinotti, A. L., et al. 2010, *A&A*, 510, A92
- McKernan, B., Yaqoob, T., & Reynolds, C. S. 2007, *MNRAS*, 379, 1359
- Misawa, T., Charlton, J. C., Eracleous, M., Ganguly, R., Tytler, D., Kirkman, D., Suzuki, N., & Lubin, D. 2007, *ApJS*, 171, 1
- Netzer, H., et al. 2003, *ApJ*, 599, 933
- Netzer, H. 2004, *ApJ*, 604, 551
- Nicastro, F., Fiore, F., Perola, G. C., & Elvis, M. 1999, *ApJ*, 512, 184
- Nicastro, F., Krongold, Y., & Elvis, M. 2008, *Memorie della Societa Astronomica Italiana*, 79, 1178
- Noble, M. S., Ji, L., Young, A., & Lee, J. C. 2009, *Astronomical Society of the Pacific Conference Series*, 411, 301
- Nucita, A. A., Guainazzi, M., Longinotti, A. L., Santos-Lleo, M., Maruccia, Y., & Bianchi, S. 2010, arXiv:1003.1285
- Penton, S. V., Stocke, J. T., & Shull, J. M. 2000, *ApJS*, 130, 121
- Piconcelli, E., Jimenez-Bailón, E., Guainazzi, M., Scharrel, N., Rodríguez-Pascual, P. M., & Santos-Lleo, M. 2005, *A&A*, 432, 15
- Proga, D., & Kallman, T. R. 2004, *ApJ*, 616, 688

- Reynolds, C. S. 1997, MNRAS, 286, 513
Risaliti, G., & Elvis, M. 2009, arXiv:0911.0958
Sako, M., et al. 2001, A&A, 365, L168
Shu, X. W., Yaqoob, T., & Wang, J. X. 2010, ApJS, 187, 581
Shull, J. M., & van Steenberg, M. 1982, ApJS, 48, 95
Shinozaki, K., Miyaji, T., Ishisaki, Y., Ueda, Y., & Ogasaka, Y. 2006, AJ, 131, 2843
Spergel D. N., et al., 2007, ApJS, 170, 377
Steenbrugge, K. C., et al. 2005, A&A, 434, 569
Steenbrugge, K. C., Fenovčík, M., Kaastra, J. S., Costantini, E., & Verbunt, F. 2009, A&A, 496, 107
Stoll, R., Mathur, S., Krongold, Y., & Nicastro, F. 2009, arXiv:0903.5310
Turner, T. J., George, I. M., Kallman, T., Yaqoob, T., & Zycki, P. T. 1996, ApJ, 472, 571
Vestergaard, M., & Peterson, B. M. 2006, ApJ, 641, 689
Wilms, J., Allen, A., & McCray, R. 2000, ApJ, 542, 914
Wood, K. S., et al. 1984, ApJS, 56, 507
Yaqoob, T., McKernan, B., Kraemer, S. B., Crenshaw, D. M., Gabel, J. R., George, I. M., & Turner, T. J. 2003, ApJ, 582, 105




Figure 1. Light curves from *Chandra* observations of Mrk 290. Each panel corresponds to an observation in Table 1. The upper half of each panel shows the hard X-ray (2-10 Å) count rate and the lower half shows the softness ratio, defined as the count rate in the 15-25 Å band divided by that in the 2-10 Å. The dashed lines serve as variability references. The count rate and softness ratio are lowest in observation 4399, while the other three observations are consistent with each other.




Figure 2. Combined MEG and HEG 1st order spectra compared to the best-fitting power-law plus black body continuum model. Residuals are given in the bottom two panels for the high-state (black line) and for the mid-state (green line). The data were binned to 0.01 Å. Possible spectral features are marked. In the residuals, C denotes for the Cash Statistic, and $\sqrt{\Delta C}$ is a proxy for χ because ΔC is distributed as χ^2 .






Figure 3. Comparison of the *XMM-Newton* *pn* spectrum to the best-fitting power-law plus black body model in the 0.5-10 keV band. Iron K α emission and an absorbing feature from 0.7 to 0.9 keV can be seen clearly in the ratio of the data to the model. The absorption is due to the warm absorber.




Figure 4. Comparison of the RGS spectrum to best-fitting power-law plus black body model from *pn* spectral fit. The Fe UTA is clearly observed in the 16.0 - 17.5 Å range, along with other features that are marked.




Figure 5. The spectral energy distributions of high-, mid- and low-states used for photoionization modeling.




Figure 6. The photoionization models for coarsely binned HETGS spectra. The upper panel shows the high-state spectrum and the lower panel shows the mid-state spectrum. In the residuals, the green lines indicate the model with one WA component, while the red lines represent the model with two WA components. The difference between the two models is primarily found at the Fe UTA feature. The residuals around 23 $\text{\AA}$ are partly due to the O VII(f) emission line.




Figure 7. The photoionization model for the low-state RGS spectrum. The lines labeled in black are at $z = 0$ and are due to Galactic gas. The O VII(f) emission line is significant.




Figure 8. Three WA components are used to model the high-state spectrum of Mrk 290, shown on a scale where 1 is the incident continuum level. The three lines of Fe XXI-XXIII marked in black indicate the highest ionized component, while the broad Fe UTA results from the lowest ionization component (in green).




Figure 9. The best fit model for high-state spectra in the 1.5-25 Å band, which consists of three WA components. Four fitted emission lines are marked in purple. The upper part and lower part of each panel show the count spectra and the flux spectra, respectively.




Figure 9. Continued.




Figure 10. The 6.7-7.5 Å regions of mid- and high-state spectra are shown in the left and right panel, respectively. The Si VII/Si VIII line at 7.23 Å is significantly stronger in the mid-state than in the high state, indicating that the ionization state of the WA responded to the continuum changes.




Figure 11. Thermal stability curve for a low-density gas illuminated by the high-state continuum of Mrk 290. The quantity $\Xi = \xi/4\pi ckT$ is the pressure ionization parameter. The black sections show the locations of the three warm absorbers, with the corresponding uncertainties on the ionization parameters. The similar value of Ξ indicates that the three components have roughly the same gas pressure.

Table 1. *Chandra* HETGS observation log

ID	Start Time	Exposure time
4399	2003.06.29 01:38:04	85.1 ks
3567	2003.07.15 01:18:47	55.1 ks
4441	2003.07.15 22:54:28	60.9 ks
4442	2003.07.17 11:18:15	50.2 ks

Table 2. *XMM-Newton* observation log

ID	Start Time	Filtered time
0400360201	2006.04.30 16:47:58	20 ks
0400360301	2006.05.02 15:21:00	7.5 ks
0400360601	2006.05.04 14:45:53	15 ks
0400360801	2006.05.06 14:39:11	14 ks

Table 3. Continuum parameters

Date	State	^a Γ	^b kT (eV)	Flux _{2–10keV} (10^{-11} erg s $^{-1}$ cm $^{-2}$)	L _{2–10keV} (10^{43} erg s $^{-1}$)	IDs and notes
2003.06.29	high-	1.86 ± 0.02	81 ± 6	1.71	3.13	3567,4441,4442
2003.07.15	mid-	1.55 ± 0.02	69 ± 9	1.20	2.19	4399
2006.04.30	low-	1.58 ± 0.01	87 ± 1	0.88	1.61	EPIC <i>pn</i> , RGS

^aThe photon index of the power-law component. ^bThe temperature of the black-body component.

Table 4. Absorption lines from the high resolution spectra with Poisson probability greater than 0.995

λ_{obs} (Å)	Flux ($10^{-6} \text{ ph cm}^{-2} \text{ s}^{-1}$)	Outflow Velocity (km s^{-1})	EWs (mÅ)	Poisson Probability	Ion Name & λ_{rest} (Å)
High-state spectrum					
4.202 ± 0.006	3.7 ± 1.8	790 ± 420	7 ± 4	0.9990	S xv (4.088)
6.360 ± 0.002	2.6 ± 0.5	530 ± 90	5 ± 1	0.9995	Si xiv Ly α (6.183)
6.844 ± 0.003	2.5 ± 0.5	270 ± 110	5 ± 1	0.9993	Si xiii(r) (6.648)
7.133 ± 0.001	2.3 ± 0.5	380 ± 60	5 ± 1	1.0000	Si ix (6.923,6.939)
8.664 ± 0.003	3.8 ± 1.4	440 ± 120	8 ± 3	1.0000	Mg xii Ly α (8.421)
9.433 ± 0.002	3.4 ± 1.0	480 ± 80	8 ± 2	0.9975	Mg xi (9.169)
9.979 ± 0.005	3.9 ± 1.4	520 ± 170	10 ± 3	0.9996	Ne x Ly γ (9.708)
10.931 ± 0.008	3.8 ± 1.8	630 ± 230	8 ± 4	0.9969	Fe xix (10.630)
12.109 ± 0.005	4.2 ± 2.0	470 ± 110	9 ± 4	0.9988	Fe xxii (11.770)
12.482 ± 0.003	10.0 ± 2.4	520 ± 70	21 ± 7	1.0000	Ne x Ly α (12.134)
12.632 ± 0.007	6.8 ± 2.8	620 ± 170	17 ± 7	1.0000	Fe xxii (12.284)
13.205 ± 0.010	9.5 ± 4.6	450 ± 240	23 ± 11	0.9999	Fe xx (12.824,12.846,12.864)
13.841 ± 0.010	21.0 ± 7.1	330 ± 220	45 ± 15	1.0000	Ne ix (13.447), Fe xix (13.423,13.375)
13.900 ± 0.001	8.8 ± 2.0	400 ± 30	19 ± 4	1.0000	Fe xix (13.497,13.518)
13.902 ± 0.008	8.9 ± 5.5	350 ± 180	21 ± 13	1.0000	Fe xxi (13.507)
14.613 ± 0.006	10.1 ± 4.6	570 ± 130	23 ± 11	1.0000	Fe xviii (14.208)
14.950 ± 0.015	5.1 ± 4.2	530 ± 300	12 ± 10	0.9998	Fe xviii (14.534)
15.446 ± 0.006	7.4 ± 4.7	490 ± 120	16 ± 10	1.0000	Fe xvii (15.014)
15.621 ± 0.015	12.1 ± 6.3	330 ± 300	24 ± 12	0.9999	O viii Ly γ (15.176)
16.472 ± 0.017	8.5 ± 8.2	380 ± 320	17 ± 16	0.9950	O viii Ly β (16.006)
16.841 ± 0.006	9.2 ± 3.6	310 ± 110	18 ± 8	1.0000	Fe x (16.360)
18.942 ± 0.006	14.5 ± 4.9	-400 ± 90	24 ± 8	0.9990	*O viii Ly α (18.967)
19.164 ± 0.002	23.4 ± 6.2	460 ± 40	35 ± 9	0.9994	O vii (18.627)
19.519 ± 0.010	18.4 ± 9.3	390 ± 170	27 ± 14	0.9950	O viii Ly α (18.967)
Mid-state spectrum					
6.357 ± 0.005	2.1 ± 1.0	660 ± 220	8 ± 4	0.9983	Si xiv Ly α (6.183)
7.229 ± 0.002	2.6 ± 0.7	-	12 ± 3	1.0000	Si viii (6.9990) or Si vii (7.063)
8.665 ± 0.005	2.3 ± 0.7	430 ± 170	11 ± 3	1.0000	Mg xii Ly α (8.421)
10.940 ± 0.005	2.6 ± 1.3	370 ± 140	13 ± 6	0.9987	Fe xix (10.630)
11.348 ± 0.006	2.7 ± 1.2	440 ± 170	14 ± 6	0.9978	Fe xix (11.029)
12.481 ± 0.003	8.6 ± 2.5	540 ± 80	21 ± 6	0.9958	Ne x Ly α (12.134)
13.205 ± 0.010	9.5 ± 4.5	450 ± 250	23 ± 11	0.9986	Fe xx (12.824,12.846,12.864)
15.624 ± 0.003	8.7 ± 3.1	260 ± 50	42 ± 15	0.9959	O viii Ly γ (15.176)
Low-state spectrum					
12.396 ± 0.081	8.7 ± 8.0	2640 ± 2000	62 ± 57	0.9984	Ne x Ly α (12.134)
13.823 ± 0.023	11.8 ± 3.3	1400 ± 520	73 ± 20	1.0000	Ne ix (13.447), Fe xix (13.497,13.518)
14.550 ± 0.018	14.0 ± 3.1	1890 ± 380	83 ± 18	1.0000	Fe xviii (14.208)
21.568 ± 0.056	22.1 ± 11.8	-490 ± 780	100 ± 41	0.9988	*O vii(r) (21.602)
24.833 ± 0.023	13.8 ± 6.6	-100 ± 280	52 ± 25	0.9981	*N vi (24.898), *N vii Ly α (24.785)
25.492 ± 0.029	11.6 ± 6.6	520 ± 350	44 ± 25	0.9977	N vii (24.779,24.785)

* Galactic absorption lines, the velocities of which are relative to the heliocentric system.

Table 5. Emission lines in the high- and low-state spectra

Ion	λ_{rest} (Å)	λ_{obs} (Å)	flux (10^{-6} ph cm $^{-2}$ s $^{-1}$)	Outflow Velocity (km s $^{-1}$)
High-state spectrum				
Fe K α	1.938	1.995 ± 0.008	4.3 ± 3.3	-
Fe XXV	1.859	$1.897^{+0.012}_{-0.028}$	10.9 ± 7.9	-
Ne IX(f)	13.669	14.115 ± 0.007	8.8 ± 4.6	-670 ± 150
O VII(f)	22.098	22.831 ± 0.008	54.7 ± 37.3	-830 ± 110
Low-state spectrum				
Fe K α	1.938	2.001 ± 0.021	4.8 ± 1.7	-
O VII(f)	22.098	22.818 ± 0.026	29.0 ± 16.5	-660 ± 350

Table 6. Best fit parameters for one WA component

State	Log ξ (erg s cm $^{-1}$)	N_H (10^{21} cm $^{-2}$)	^{a} Velocity (km s $^{-1}$)	C stat/d.o.f.	$^b \Delta C$
high-	2.43 ± 0.02	4.9 ± 0.5	450 ± 30	2635/2344	431
mid-	2.41 ± 0.07	4.8 ± 1.3	450 ± 60	2071/1843	92

^a The outflow velocity of WA is relative to the systematic velocity of Mrk 290.^b Change in the C statistics between models with and without a warm absorber.**Table 7.** The two phase WA parameters for three states.

State	$^a \Gamma$	$^b kT$ (eV)	$L_{1-1000 \text{ Ryd}}$ (10^{43} erg s $^{-1}$)		Log ξ (erg s cm $^{-1}$)	N_H (10^{21} cm $^{-2}$)	^{c} Velocity (km s $^{-1}$)	C stat/d.o.f.
high-	1.91 ± 0.01	119 ± 3	13.7	WA1	1.62 ± 0.15	0.54 ± 0.18	540 ± 150	414/306
				WA2	2.45 ± 0.04	5.33 ± 1.17	$^d 450$	
^{e} mid-	1.64 ± 0.02	95^{+6}_{-1}	10.0	WA1	1.52 ± 0.55	0.54	540	172/152
				WA2	2.37 ± 0.08	5.33	450	
low-	1.69 ± 0.01	105 ± 2	7.2	WA1	1.57 ± 0.15	1.12 ± 0.18	1260 ± 160	225/183
				WA2	$2.50^{+0.31}_{-0.08}$	$3.16^{+0.35}_{-1.48}$	1260 ± 250	

^a The photon index of the power-law component in the intrinsic spectra.^b The temperature of the black-body component in the intrinsic spectra.^c The outflow velocity of WA is relative to the systematic velocity of Mrk 290.^d The outflow velocity is fixed to the value in Table 6.^e The column density and the outflow velocity of WAs are fixed to the values obtained from the high-state spectral fits.**Table 8.** Best fit parameters for the three WAs in the high-state spectrum

	Log ξ (erg s cm $^{-1}$)	N_H (10^{21} cm $^{-2}$)	^{a} Velocity (km s $^{-1}$)
^{b} WA1	1.62	0.54	540
WA2	2.42 ± 0.04	4.06 ± 0.62	450 ± 30
WA3	3.20 ± 0.14	3.49 ± 1.47	390 ± 60

^a The outflow velocity of WA is relative to the systematic velocity of Mrk 290.^b The parameters of this lowest ionization component is fixed to the values in Table 7.

